

SPECTRAL INSTABILITY OF THE PEAKED PERIODIC WAVE IN THE REDUCED OSTROVSKY EQUATIONS

ANNA GEYER AND DMITRY E. PELINOVSKY

(Communicated by Catherine Sulem)

ABSTRACT. We show that the peaked periodic traveling wave of the reduced Ostrovsky equations with quadratic and cubic nonlinearity is spectrally unstable in the space of square integrable periodic functions with zero mean and the same period. We discover that the spectrum of a linearized operator at the peaked periodic wave completely covers a closed vertical strip of the complex plane. In order to obtain this instability, we prove an abstract result on spectra of operators under compact perturbations. This justifies the truncation of the linearized operator at the peaked periodic wave to its differential part for which the spectrum is then computed explicitly.

1. INTRODUCTION

The Ostrovsky equation with quadratic nonlinearity was originally derived by L.A. Ostrovsky [18] to model small-amplitude long waves in a rotating fluid of finite depth. The same approximation was extended to internal gravity waves, in which case the underlying equation includes the cubic nonlinearity and is referred to as the modified Ostrovsky equation [9, 11, 19]. When the high-frequency dispersion is neglected, the reduced Ostrovsky equation can be written in the form

$$(1.1) \quad u_t + uu_x = \partial_x^{-1}u,$$

whereas the reduced modified Ostrovsky equation takes the form

$$(1.2) \quad u_t + u^2u_x = \partial_x^{-1}u.$$

For both equations (1.1) and (1.2), periodic waves of the normalized period 2π are considered in the Sobolev space of 2π -periodic functions denoted by $H_{\text{per}}^s \equiv H_{\text{per}}^s(-\pi, \pi)$ for some $s \geq 0$. The subspace of H_{per}^s for 2π -periodic functions with zero mean is denoted by \dot{H}_{per}^s . The operator $\partial_x^{-1} : \dot{H}_{\text{per}}^s \rightarrow \dot{H}_{\text{per}}^{s+1}$ denotes the anti-derivative with zero mean.

Local well-posedness of the Cauchy problem for the reduced Ostrovsky equations (1.1) and (1.2) can be shown in \dot{H}_{per}^s with $s > \frac{3}{2}$ [16, 21]. For sufficiently large initial data, the local solutions break in finite time, similar to the inviscid Burgers equation

Received by the editors July 18, 2019, and, in revised form, July 31, 2019, and October 23, 2019.

2010 *Mathematics Subject Classification.* Primary 35B35, 35Q35.

Key words and phrases. Peaked periodic wave, reduced Ostrovsky equation, spectral instability.

The second author acknowledges financial support from the state task program in the sphere of scientific activity of the Ministry of Science and Higher Education of the Russian Federation (Task No. FSWE-2020-0007) and from the grant of the president of the Russian Federation for the leading scientific schools (grant No. NSH-2485.2020.5).

[5, 10, 16]. For sufficiently small initial data, the local solutions are continued for all times [12].

Traveling wave solutions of the reduced Ostrovsky equations are of the form $u(x, t) = U(x - ct)$, where $z = x - ct$ is the traveling wave coordinate and c is the wave speed. The wave profile $U(z)$ satisfies the integral-differential equation in the form

$$(1.3) \quad \begin{cases} [c - U(z)^p] U'(z) + (\partial_z^{-1} U)(z) = 0, & \forall z \in (-\pi, \pi) \text{ with } U(z) \neq c, \\ U(-\pi) = U(\pi), \quad \int_{-\pi}^{\pi} U(z) dz = 0, \end{cases}$$

where $p = 1$ for (1.1) and $p = 2$ for (1.2).

Smooth solutions to the boundary-value problem (1.3) exist for $c \in (1, c_*)$, where c_* is uniquely defined; see [7] (and [1] for a generalization). For $c \in (1, c_*)$ smooth solutions satisfy $U(z) < c$ for every $z \in [-\pi, \pi]$, and the boundary-value problem (1.3) can be equivalently rewritten in the differential form

$$(1.4) \quad \begin{cases} \frac{d}{dz} [(c - U(z)^p) \frac{dU}{dz}] + U(z) = 0, & \text{for every } z \in [-\pi, \pi], \\ U(-\pi) = U(\pi), \quad U'(-\pi) = U'(\pi). \end{cases}$$

At $c = c_*$ solutions to the boundary-value problem (1.3) are peaked at the points $z = \pm\pi$ where $U(\pm\pi) = c_*$. Uniqueness and Lipschitz continuity of the peaked solutions to the boundary-value problem (1.3) were proven in [8] for $p = 1$ (see [1, 4] for a generalization). We denote this unique (up to translation) peaked solution by $U_*(z)$.

For $p = 1$, the peaked wave $U_*(z)$ exists at the wave speed $c_* = \frac{\pi^2}{9}$ and is given by

$$(1.5) \quad U_*(z) = \frac{1}{18}(3z^2 - \pi^2), \quad \text{for } z \in [-\pi, \pi],$$

periodically continued beyond $[-\pi, \pi]$. It was already obtained in the original paper [18]. For $p = 2$, the peaked wave $U_*(z)$ exists at the wave speed $c_* = \frac{\pi^2}{8}$ and is given by

$$(1.6) \quad U_*(z) = \frac{1}{\sqrt{2}} \left(|z| - \frac{\pi}{2} \right), \quad \text{for } z \in [-\pi, \pi],$$

periodically continued beyond $[-\pi, \pi]$; see [17]. In both cases, $U_* \in \dot{H}_{\text{per}}^s$ for $s < \frac{3}{2}$ with a finite jump discontinuity of the first derivative at $z = \pm\pi$ for (1.5) and at $z = 0, \pm\pi$ for (1.6).

Smooth periodic waves of the quasi-linear differential equation in (1.4) can be obtained equivalently from a semi-linear differential equation by means of the following change of coordinates [6, 13, 14]:

$$(1.7) \quad U(z) = \mathbf{u}(\xi), \quad z = \int_0^\xi [c - \mathbf{u}(s)^p] ds.$$

The smooth periodic waves with profile \mathbf{u} satisfy the differential equation

$$(1.8) \quad \frac{d^2 \mathbf{u}}{d\xi^2} + [c - \mathbf{u}(\xi)^p] \mathbf{u}(\xi) = 0.$$

Although all periodic solutions of differential equation (1.8) are smooth, the coordinate transformation (1.7) fails to be invertible if $\mathbf{u}(\xi) = c$ for some ξ . Singularities in the coordinate transformation are related to the appearance of the peaked solutions in the boundary-value problem (1.3).

Spectral stability of smooth periodic waves with respect to perturbations of the same period was proven both for (1.1) and (1.2) in [7, 14]. The analysis of [7] relies on the standard variational formulation of the periodic waves as critical points of energy subject to fixed momentum. The analysis of [14] relies on the coordinate transformation (1.7), which reduces the spectral stability problem of the form $\partial_x Lv = \lambda v$ with the self-adjoint operator $L = c - U^p + \partial_z^{-2}$ to the spectral problem of the form $M\mathbf{v} = \lambda \partial_\xi \mathbf{v}$ with the self-adjoint operator $M = c - \mathbf{u}^p + \partial_\xi^2$. The spectral problem $M\mathbf{v} = \lambda \partial_\xi \mathbf{v}$ has been studied before in [22] (see also [15] for a generalization). Orbital stability of smooth periodic waves with respect to perturbations of any period multiple to the wave period was proven in [6] by using higher-order conserved quantities of the reduced Ostrovsky equations (1.1) and (1.2).

The peaked periodic waves are, informally speaking, located at the boundary between global and breaking solutions in the reduced Ostrovsky equations. If the initial data u_0 is smooth, it was shown that global solutions of (1.1) exist if $m_0(x) := 1 - 3u_0''(x) > 0$ for every x and wave breaking occurs if $m_0(x)$ is sign-indefinite [10, 12], whereas global solutions of (1.2) exist if $m_0(x) := 1 - \sqrt{2}|u_0'(x)| > 0$ for every x and wave breaking occurs if $m_0(x)$ is sign-indefinite [5]. Substituting U_* instead of u_0 yields $m_0(x) = 0$ almost everywhere except at the peaks. Thus, it is natural to expect that the peaked periodic waves are unstable in the time evolution of the reduced Ostrovsky equations.

In [8] we proved that the unique peaked solution (1.5) of the reduced Ostrovsky equation (1.1) is linearly unstable with respect to square integrable perturbations with zero mean and the same period. This was done by obtaining sharp bounds on the exponential growth of the L^2 norm of the perturbations in the linearized time-evolution problem $v_t = \partial_z Lv$. No claims regarding the spectral instability of the peaked periodic wave were made in [8]. In [14], explicit solutions of the spectral stability problem $M\mathbf{v} = \lambda \partial_\xi \mathbf{v}$ were constructed, but since this construction violated the periodic boundary conditions on the perturbation term, it did not provide an answer to the spectral stability question.

The main goal of this paper is to show that *the peaked periodic wave U_* is spectrally unstable with respect to square integrable perturbations with zero mean and the same period*. We achieve this for both versions of the reduced Ostrovsky equations (1.1) and (1.2) with the peaked periodic waves U_* given in (1.5) and (1.6), respectively. We discover *an unusual instability of the peaked periodic wave*: the spectrum of the linearized operator $A = \partial_z L$ in the space of 2π -periodic mean-zero functions completely covers a closed vertical strip of the complex plane, as depicted in Figure 1 for the reduced Ostrovsky equation (1.1). The right boundary of this vertical strip with $\operatorname{Re}(\lambda) = \frac{\pi}{6}$ coincides with the sharp growth rate of the exponentially growing perturbations obtained in [8] for the peaked wave U_* given by (1.5). The vertical strip remains invariant when the spectrum of A is defined in the space of subharmonic and localized perturbations; see Remark 4. A similar instability with the spectrum lying in a vertical strip was discovered in [20] in the context of linearization around double periodic steady state solutions of the 2D Euler equations.

Let us recall the following standard definition (see Definition 6.1.9 in [2]).




FIGURE 1. The spectrum of the linearized operator A at the peaked periodic wave U_* given by (1.5) completely covers a closed vertical strip in the complex plane with zero being the only eigenvalue. This shows that the peaked wave is spectrally unstable with respect to coproduct perturbations.

Definition 1. Let A be a linear operator on a Banach space X with $\text{dom}(A) \subset X$. The complex plane \mathbb{C} is decomposed into the following two sets:

- (1) The resolvent set

$$\rho(A) = \{ \lambda \in \mathbb{C} : \ker(A - \lambda I) = \{0\}, \text{ran}(A - \lambda I) = X, (A - \lambda I)^{-1} : X \rightarrow X \text{ is bounded} \}.$$

- (2) The spectrum

$$\sigma(A) = \mathbb{C} \setminus \rho(A),$$

which is further decomposed into the following three disjoint sets:

- (a) the point spectrum

$$\sigma_p(A) = \{ \lambda \in \sigma(A) : \ker(A - \lambda I) \neq \{0\} \},$$

- (b) the residual spectrum

$$\sigma_r(A) = \{ \lambda \in \sigma(A) : \ker(A - \lambda I) = \{0\}, \text{ran}(A - \lambda I) \neq X \},$$

- (c) the continuous spectrum

$$\sigma_c(A) = \{ \lambda \in \sigma(A) : \ker(A - \lambda I) = \{0\}, \text{ran}(A - \lambda I) = X, (A - \lambda I)^{-1} : X \rightarrow X \text{ is unbounded} \}.$$

In order to prove the spectral instability of the peaked periodic waves, we proceed as follows. We first show that the point spectrum of the linearized operator A consists of only the zero eigenvalue; see Lemma 1. We then observe that A is the sum of the linearization A_0 of the quasi-linear part of the equation and a nonlocal term, which we may view as a compact perturbation K . The truncated spectral problem for A_0 is then transformed to a problem on the line by a change of coordinates in Lemma 2. This facilitates the explicit computation of the spectrum of A_0 in Lemmas 3 and 4. Finally, we justify the truncation of the linearized operator to its differential part by verifying the assumptions of the following abstract result, which is proven in the appendix.

Theorem 1. *Let $A : \text{dom}(A) \subset X \rightarrow X$ and $A_0 : \text{dom}(A_0) \subset X \rightarrow X$ be linear operators on Hilbert space X with the same domain $\text{dom}(A_0) = \text{dom}(A)$ such that $A - A_0 = K$ is a compact operator in X . Assume that the intersections $\sigma_p(A) \cap \rho(A_0)$ and $\sigma_p(A_0) \cap \rho(A)$ are empty. Then, $\sigma(A) = \sigma(A_0)$.*

The proof of nonlinear instability of the peaked periodic waves is still open for the reduced Ostrovsky equations (1.1) and (1.2). One of the main obstacles for nonlinear stability analysis is the lack of well-posedness results for initial data in \dot{H}_{per}^s with $s < \frac{3}{2}$, which would include the peaked periodic waves U_* given by (1.5) and (1.6). Another obstacle is the discrepancy between the domain of the linearized operator $A = \partial_z L = \partial_z(c_* - U_*^p) + \partial_z^{-1}$ in \dot{L}_{per}^2 and the Sobolev space \dot{H}_{per}^1 : while the former allows finite jumps of perturbations at the peaks, the latter requires continuity of perturbations across the peaks; see Remark 3.

The paper is organized as follows. The linearized operator A is studied in Section 2, where the main results for the peaked periodic waves U_* given by (1.5) and (1.6) are formulated. The proofs of the main results are contained in Sections 3 and 4. The appendix contains the proof of Theorem 1.

2. MAIN RESULTS

Linearizing (1.1) or (1.2) about the peaked traveling wave $U_*(x - c_*t)$ with the perturbation $v(t, x - c_*t)$ yields an evolution problem of the form

$$(2.1) \quad v_t = Av,$$

where the operator $A : \text{dom}(A) \subset \dot{L}_{\text{per}}^2 \rightarrow \dot{L}_{\text{per}}^2$ is defined by

$$(2.2) \quad (Av)(z) := \partial_z [(c_* - U_*(z)^p)v(z)] + \partial_z^{-1}v(z), \quad z \in (-\pi, \pi),$$

with maximal domain

$$(2.3) \quad \text{dom}(A) = \left\{ v \in \dot{L}_{\text{per}}^2 : \partial_z [(c_* - U_*^p)v] \in \dot{L}_{\text{per}}^2 \right\},$$

where either $p = 1$ for (1.1) or $p = 2$ for (1.2).

The linearized operator (2.2) can be written as $A = A_0 + K$, where the truncated operator $A_0 : \text{dom}(A_0) \subset \dot{L}_{\text{per}}^2 \rightarrow \dot{L}_{\text{per}}^2$ is defined by

$$(2.4) \quad (A_0v)(z) := \partial_z [(c_* - U_*(z)^p)v(z)], \quad z \in (-\pi, \pi),$$

with the same domain $\text{dom}(A_0) = \text{dom}(A)$ and $K := \partial_z^{-1}$ is a compact (Hilbert-Schmidt) operator in \dot{L}_{per}^2 with spectrum $\sigma(K) = \{in^{-1}, n \in \mathbb{Z} \setminus \{0\}\}$.

By using Definition 1, we introduce the following notion of spectral stability for the traveling wave U_* .

Definition 2. The traveling wave U_* is said to be spectrally stable if $\sigma(A) \subset i\mathbb{R}$. Otherwise, it is said to be spectrally unstable.

The following two theorems present the main results of this paper.

Theorem 2. *Consider the operator A given by (2.2) on \dot{L}_{per}^2 with $\text{dom}(A)$ given by (2.3) for $p = 1$ and U_* as in (1.5). Then,*

$$(2.5) \quad \sigma(A) = \left\{ \lambda \in \mathbb{C} : -\frac{\pi}{6} \leq \text{Re}(\lambda) \leq \frac{\pi}{6} \right\}.$$

Consequently, the peaked wave U_ is spectrally unstable in the reduced Ostrovsky equation (1.1).*

Theorem 3. Consider the operator A given by (2.2) on \dot{L}_{per}^2 with $\text{dom}(A)$ given by (2.3) for $p = 2$ and U_* in (1.6). Then,

$$(2.6) \quad \sigma(A) = \left\{ \lambda \in \mathbb{C} : -\frac{\pi}{4} \leq \text{Re}(\lambda) \leq \frac{\pi}{4} \right\}.$$

Consequently, the peaked wave U_* is spectrally unstable in the reduced modified Ostrovsky equation (1.2).

Remark 1. One can find the explicit eigenvector for $0 \in \sigma(A)$ thanks to the translational symmetry implying $AU'_* = 0$, where $U'_* \in \text{dom}(A) \subset \dot{L}_{\text{per}}^2$. Therefore, $0 \in \sigma_p(A)$. We show in Lemmas 1 and 5 that $\sigma_p(A) = \{0\}$.

Remark 2. We are not able to distinguish between residual and continuous spectrums in $\sigma(A) \setminus \{0\}$. This is because we truncate the operator A to an operator A_0 with the same domain and use the result of Theorem 1. For the operator A_0 we prove in Lemmas 2, 3, 4, and 6 that $\sigma_p(A_0)$ is empty and $\sigma_r(A_0)$ is the open vertical strip in (2.5) and (2.6), whereas $\sigma_c(A_0)$ is the boundary of that vertical strip.

Remark 3. The Sobolev space \dot{H}_{per}^1 is continuously embedded into $\text{dom}(A)$ in the sense that there exists $C > 0$ such that for every $f \in \dot{H}_{\text{per}}^1$, we have $\partial_z [(c_* - U_*^p)f] \in \dot{L}_{\text{per}}^2$ with the bound

$$\|\partial_z [(c_* - U_*^p)f]\|_{L_{\text{per}}^2} \leq C \|f\|_{\dot{H}_{\text{per}}^1}.$$

However, \dot{H}_{per}^1 is not equivalent to $\text{dom}(A)$ because piecewise continuous functions with finite jump discontinuities at the points z where $c_* - U_*^p(z)$ vanishes belong to $\text{dom}(A)$ but do not belong to \dot{H}_{per}^1 . For example, the eigenvector $U'_* \in \text{dom}(A)$ for $0 \in \sigma_p(A)$ does not belong to \dot{H}_{per}^1 .

3. PROOF OF THEOREM 2

For the peaked periodic wave U_* in (1.5) in the case $p = 1$, we write explicitly

$$(3.1) \quad c_* - U_*(z) = \frac{1}{6} [\pi^2 - z^2], \quad z \in [-\pi, \pi].$$

The eigenvector for $0 \in \sigma_p(A)$ is given by

$$(3.2) \quad U'_*(z) = \frac{1}{3}z, \quad z \in (-\pi, \pi).$$

The proof of Theorem 2 can be divided into four steps.

Step 1 (Point spectrum of A). If $\lambda \in \sigma_p(A)$, then there exists $f \in \text{dom}(A)$, $f \neq 0$, such that $Af = \lambda f$. It follows from Remark 1 that $0 \in \sigma_p(A)$ with the eigenvector U'_* in (3.2). The following result shows that no other eigenvalues of $\sigma_p(A)$ exist.

Lemma 1. $\sigma_p(A) = \{0\}$.

Proof. First we note that if $f \in \text{dom}(A)$, then $f \in H^1(-\pi, \pi)$ so that $f \in C^0(-\pi, \pi)$ by Sobolev embedding. Bootstrapping arguments for $Af = \lambda f$ immediately yield that $f \in C^\infty(-\pi, \pi)$; hence the spectral problem $Af = \lambda f$ for $f \in \text{dom}(A)$ can be differentiated once in z to yield the second-order differential equation

$$(3.3) \quad (\pi^2 - z^2)f''(z) - 4zf'(z) + 4f(z) = 6\lambda f'(z), \quad z \in (-\pi, \pi).$$

One solution is available in closed form: $f_1(z) = 2z + 3\lambda$. In order to obtain the second linearly independent solution, we write $f_2(z) = (2z + 3\lambda)g(z)$ and derive the following equation for g :

$$(3.4) \quad (\pi^2 - z^2)(2z + 3\lambda)g''(z) + 2[2(\pi^2 - z^2) - (2z + 3\lambda)^2]g'(z) = 0, \quad z \in (-\pi, \pi).$$

This equation can be integrated once to obtain

$$(3.5) \quad g'(z) = \frac{g_0}{(\pi^2 - z^2)^2(2z + 3\lambda)^2} \left(\frac{\pi + z}{\pi - z} \right)^{\frac{3\lambda}{\pi}}, \quad z \in (-\pi, \pi),$$

where g_0 is a constant of integration. Computing the limits $z \rightarrow \pm\pi$ shows that if $\pm 2\pi + 3\lambda \neq 0$, then

$$g'(z) \sim \begin{cases} (\pi - z)^{-\frac{3\lambda}{\pi} - 2}, & z \rightarrow \pi, \\ (\pi + z)^{\frac{3\lambda}{\pi} - 2}, & z \rightarrow -\pi. \end{cases}$$

This sharp asymptotical behavior shows that $(\pi^2 - z^2)g'(z) \notin L^2(-\pi, \pi)$, even if $g \in L^2(-\pi, \pi)$. Therefore, for every $\lambda \in \mathbb{C}$ with $\pm 2\pi + 3\lambda \neq 0$, the second solution $f_2(z)$ does not belong to $\text{dom}(A) \subset \dot{L}_{\text{per}}^2$ because of the divergences as $z \rightarrow \pm\pi$. For $\pm 2\pi + 3\lambda = 0$, the explicit expression (3.5) yields

$$g'(z) = \frac{g_0}{4(\pi^2 - z^2)^2(\pi \pm z)^2}, \quad z \in (-\pi, \pi),$$

which still implies that f_2 does not belong to $\text{dom}(A) \subset \dot{L}_{\text{per}}^2$. Hence, for every $\lambda \in \mathbb{C}$, if $f \in \text{dom}(A) \subset \dot{L}_{\text{per}}^2$ is a solution to $Af = \lambda f$, then f is proportional to $f_1(z) = 2z + 3\lambda$ only. The zero-mass constraint $\int_{-\pi}^{\pi} f_1(z) dz = 0$ required for $f_1 \in \dot{L}_{\text{per}}^2$ yields $\lambda = 0$, so that $f_1(z) = 2z = 6U'_*(z)$ given by (3.2). No other $\lambda \in \mathbb{C}$ such that a nonzero solution f of $Af = \lambda f$ belongs to $\text{dom}(A) \subset \dot{L}_{\text{per}}^2$ exists. \square

Step 2 (Truncation of \mathbf{A}). By using (3.1), A_0 in (2.4) is rewritten in the explicit form

$$(3.6) \quad (A_0 v)(z) = \frac{1}{6} \partial_z [(\pi^2 - z^2)v(z)], \quad z \in (-\pi, \pi).$$

Inserting the expression (3.1) in the transformation formula (1.7) for $p = 1$ yields

$$(3.7) \quad \frac{dz}{d\xi} = \frac{1}{6}(\pi^2 - z^2),$$

which we can solve to find that

$$(3.8) \quad z = \pi \tanh\left(\frac{\pi\xi}{6}\right),$$

where the constant of integration is defined without loss of generality from the condition that $z = 0$ at $\xi = 0$. By using the explicit transformation formula (3.8), we can rewrite the spectral problem $A_0 v = \lambda v$ in an equivalent but more convenient form.

Lemma 2. *The spectral problem $A_0 v = \lambda v$ with $A_0 : \text{dom}(A_0) \subset \dot{L}_{\text{per}}^2 \rightarrow \dot{L}_{\text{per}}^2$ given by (3.6) is equivalent to the spectral problem $B_0 w = \mu w$ with*

$$(3.9) \quad \mu = \frac{6}{\pi}\lambda,$$

where $B_0 : \text{dom}(B_0) \subset \tilde{L}^2(\mathbb{R}) \rightarrow \tilde{L}^2(\mathbb{R})$ is the linear operator given by

$$(3.10) \quad (B_0 w)(y) := \partial_y w(y) - \tanh(y)w(y), \quad y \in \mathbb{R},$$

with maximal domain

$$(3.11) \quad \text{dom}(B_0) = \left\{ w \in \tilde{L}^2(\mathbb{R}) : (\partial_y - \tanh y)w \in \tilde{L}^2(\mathbb{R}) \right\} = H^1(\mathbb{R}) \cap \tilde{L}^2(\mathbb{R}),$$

where $\tilde{L}^2(\mathbb{R})$ is the constrained L^2 space given by

$$(3.12) \quad \tilde{L}^2(\mathbb{R}) := \{w \in L^2(\mathbb{R}) : \langle w, \varphi \rangle = 0\}$$

with $\varphi(y) := \text{sech}(y)$.

Proof. We first show that $v \in L^2(-\pi, \pi)$ if and only if $w \in L^2(\mathbb{R})$. To this end, we use the substitution rule with (3.8), set $y := \frac{\pi\xi}{6}$, and write $v(z) = \cosh(y)w(y)$ to obtain that

$$\int_{-\pi}^{\pi} v^2(z) dz = \pi \int_{-\infty}^{\infty} v^2(\pi \tanh y) \text{sech}^2(y) dy = \pi \int_{-\infty}^{\infty} w^2(y) dy.$$

Similarly, the zero-mean constraint in \dot{L}^2_{per} is transformed to

$$0 = \int_{-\pi}^{\pi} v(z) dz = \pi \int_{-\infty}^{\infty} v(\pi \tanh y) \text{sech}^2(y) dy = \pi \int_{-\infty}^{\infty} w(y) \text{sech}(y) dy.$$

Therefore, $v \in \dot{L}^2_{\text{per}}$ if and only if $w \in \tilde{L}^2(\mathbb{R})$. Furthermore, we verify that

$$\partial_z [(\pi^2 - z^2)v] \in L^2(-\pi, \pi)$$

if and only if

$$\partial_y w - \tanh(y)w \in L^2(\mathbb{R}).$$

Next we note that $B_0 w \in \tilde{L}^2(\mathbb{R})$ for every $w \in H^1(\mathbb{R})$, since

$$(3.13) \quad \langle B_0 w, \varphi \rangle = \int_{\mathbb{R}} [w'(y) - \tanh(y)w(y)] \text{sech}(y) dy = \int_{\mathbb{R}} \frac{d}{dy} [w(y)\text{sech}(y)] dy = 0.$$

This implies that the constraint $\langle B_0 w, \varphi \rangle = 0$ is identically satisfied for every $w \in H^1(\mathbb{R})$. Moreover, if $w \in L^2(\mathbb{R})$ and $[\partial_y - \tanh(y)]w \in L^2(\mathbb{R})$, then $w \in H^1(\mathbb{R})$. The above arguments show that B_0 is closed in $\tilde{L}^2(\mathbb{R})$ and $\text{dom}(B_0) = H^1(\mathbb{R}) \cap \tilde{L}^2(\mathbb{R})$. Hence, the spectral problems for A_0 and B_0 are equivalent to each other, and the spectral parameters λ and μ are related by the transformation formula (3.9). \square

Step 3 (Spectrum of the truncated operator \mathbf{A}_0). In view of the equivalence of the spectral problems of A_0 and B_0 proven in Lemma 2, we proceed to study the spectrum of B_0 in $\tilde{L}^2(\mathbb{R})$. The following two lemmas characterize the spectrum of B_0 .

Lemma 3. *The point spectrum of B_0 is empty.*

Proof. Let $\mu \in \mathbb{C}$ and $w \in \ker(B_0 - \mu I)$; i.e., w satisfies the first-order differential equation

$$\frac{dw}{dy} = \mu w(y) + \tanh(y)w(y).$$

Solving this homogeneous equation yields

$$w(y) = C \cosh(y)e^{\mu y}$$

where C is arbitrary. We have $w(y) \sim e^{(\mu \pm 1)y}$ as $y \rightarrow \pm\infty$, and hence the two exponential functions decay to zero as $y \rightarrow \pm\infty$ in two disjoint sets of \mathbb{C} for μ . Hence, $w \in \text{dom}(B_0) \subset \tilde{L}^2(\mathbb{R})$ if and only if $C = 0$ for every $\mu \in \mathbb{C}$. We conclude that $w = 0$, so $\sigma_p(B_0) = \emptyset$. \square

Lemma 4. *The residual spectrum of B_0 is*

$$(3.14) \quad \sigma_r(B_0) = \{\mu \in \mathbb{C} : -1 < \text{Re}(\mu) < 1\},$$

whereas the continuous spectrum of B_0 is

$$(3.15) \quad \sigma_c(B_0) = \{\mu \in \mathbb{C} : \text{Re}(\mu) = \pm 1\}.$$

Proof. Let $f \in \tilde{L}^2(\mathbb{R})$, $\mu \in \mathbb{C}$, and consider the resolvent equation $(B_0 - \mu I)w = f$, i.e.

$$(3.16) \quad \frac{dw}{dy} - \tanh(y)w(y) - \mu w(y) = f(y).$$

Since the spectrum $\sigma(B_0)$ is invariant under translations along the imaginary axis, it suffices to study equation (3.16) for $\mu \in \mathbb{R}$; see also Theorem 3.13 in [3]. In what follows, we will study for which $\mu \in \mathbb{R}$ the resolvent equation (3.16) has a solution w in $\text{dom}(B_0)$. Note that if $\mu \neq 0$ and $w \in H^1(\mathbb{R})$ is a solution to (3.16), then the constraint $\langle f, \varphi \rangle = 0$ implies that $\langle w, \varphi \rangle = 0$, so that $w \in H^1(\mathbb{R})$ implies that $w \in \text{dom}(B_0) = H^1(\mathbb{R}) \cap \tilde{L}^2(\mathbb{R})$. On the other hand, if $\mu = 0$ and $w \in H^1(\mathbb{R})$ is a solution to (3.16), then the constraint $\langle w, \varphi \rangle = 0$ is needed to ensure that $w \in \text{dom}(B_0)$.

Solving the first-order inhomogeneous equation (3.16) by variation of parameters yields

$$(3.17) \quad w(y) = \cosh(y)e^{\mu y} \left[C + \int_0^y e^{-\mu y'} \text{sech}(y') f(y') \, dy' \right],$$

from which we infer that $w \in H_{\text{loc}}^1(\mathbb{R})$. However, we also need to consider the behavior of $w(y)$ as $y \rightarrow \pm\infty$ to ensure that $w \in \text{dom}(B_0)$.

Let us first show that the half line $I_+ := \{\mu \in \mathbb{R} : \mu > 1\}$ belongs to the resolvent set of B_0 . Since $e^{(\mu+1)y}$ diverges as $y \rightarrow +\infty$ for every $\mu \in I_+$, we define C in (3.17) by

$$(3.18) \quad C := - \int_0^\infty e^{-\mu y'} \text{sech}(y') f(y') \, dy',$$

so that the unique solution (3.17) can be rewritten as

$$(3.19) \quad w(y) = \int_{+\infty}^y e^{\mu(y-y')} \frac{\cosh(y)}{\cosh(y')} f(y') \, dy'.$$

The following two equivalent representations will be useful in the estimates below:

$$(3.20) \quad \frac{\cosh(y)}{\cosh(y')} = \frac{1 + e^{2y}}{1 + e^{2y'}} e^{y'-y}$$

$$(3.21) \quad = \frac{1 + e^{-2y}}{1 + e^{-2y'}} e^{y-y'}.$$

Let $f = f\chi_{\{y>0\}} + f\chi_{\{y<0\}}$, where χ_S is the characteristic function on the set $S \subset \mathbb{R}$, and define w_\pm by (3.19) with f replaced by $f\chi_{\{\pm y>0\}}$ so that $w = w_+ + w_-$. Using

(3.20) for $y < 0$ and (3.21) for $y > 0$, we obtain

$$\text{for } y < 0 : \quad |w_+(y)| \leq \int_0^{+\infty} e^{-(\mu-1)(y'-y)} |f(y')| dy'$$

and

$$\text{for } y > 0 : \quad |w_+(y)| \leq 2 \int_y^{+\infty} e^{-(\mu+1)(y'-y)} |f(y')| dy'.$$

By the Cauchy–Schwarz inequality for $y < 0$ and by the generalized Young’s inequality for $y > 0$, we obtain

$$\|w_+\|_{L^2(\mathbb{R}^-)} \leq \|e^{-(\mu-1)\cdot}\|_{L^2(\mathbb{R}^+)} \|f\|_{L^2(\mathbb{R}^+)} \|e^{(\mu-1)\cdot}\|_{L^2(\mathbb{R}^-)} \leq \frac{1}{2(\mu-1)} \|f\|_{L^2(\mathbb{R}^+)}$$

and

$$\|w_+\|_{L^2(\mathbb{R}^+)} \leq 2 \|e^{-(\mu+1)\cdot}\|_{L^1(\mathbb{R}^+)} \|f\|_{L^2(\mathbb{R}^+)} \leq \frac{2}{\mu+1} \|f\|_{L^2(\mathbb{R}^+)}.$$

On the other hand, $w_-(y) = 0$ for $y > 0$ and

$$y < 0 : \quad |w_-(y)| \leq 2 \int_y^0 e^{-(\mu-1)(y'-y)} |f(y')| dy',$$

where the representation (3.20) has been used. By the generalized Young’s inequality, we obtain

$$\|w_-\|_{L^2(\mathbb{R})} \leq 2 \|e^{-(\mu-1)\cdot}\|_{L^1(\mathbb{R}^+)} \|f\|_{L^2(\mathbb{R}^-)} \leq \frac{2}{\mu-1} \|f\|_{L^2(\mathbb{R}^-)}.$$

Putting these bounds together yields

$$(3.22) \quad \|w\|_{L^2(\mathbb{R})} \leq C_\mu \|f\|_{L^2(\mathbb{R})},$$

where the constant $C_\mu > 0$ depends on μ and is bounded for every $\mu > 1$. Thus, we have showed that $I_+ \in \rho(B_0)$. Similarly, one can show that $I_- := \{\mu \in \mathbb{R} : \mu < -1\}$ also belongs to the resolvent set of B_0 due to the same bound (3.22) for every $\mu \in I_-$. Hence, $I_+ \cup I_- \subseteq \rho(B_0)$. It remains to show that $[-1, 1] \subseteq \sigma(B_0)$. More precisely, we show that $\mu \in \sigma_r(B_0)$ if $\mu \in (-1, 1)$ and $\mu \in \sigma_c(B_0)$ if $\mu = \pm 1$. We use again the explicit solution $w \in H_{loc}^1(\mathbb{R})$ given in (3.17).

If $\mu \in (-1, 1)$, then the exponential functions $e^{(\mu+1)y}$ and $e^{(\mu-1)y}$ do not decay to zero as $y \rightarrow +\infty$ and $y \rightarrow -\infty$, respectively. Therefore, to ensure decay of $w(y)$ as $y \rightarrow \pm\infty$, the constant C in (3.17) would have to be defined twice:

$$(3.23) \quad C = - \int_0^{+\infty} e^{-\mu y'} \operatorname{sech}(y') f(y') dy' = \int_{-\infty}^0 e^{-\mu y'} \operatorname{sech}(y') f(y') dy'.$$

This implies that $f \in \tilde{L}^2(\mathbb{R})$ would have to satisfy an additional constraint

$$(3.24) \quad \int_{\mathbb{R}} e^{-\mu y'} \operatorname{sech}(y') f(y') dy' = 0,$$

which is different from $\langle f, \varphi \rangle = 0$ if $\mu \neq 0$. Fix $\mu \in \mathbb{R}$ such that $\mu \in (-1, 1)$ and $\mu \neq 0$. If $f \in \tilde{L}^2(\mathbb{R})$ satisfies (3.24), then there exists a solution $w \in \operatorname{dom}(B_0)$ to the resolvent equation (3.16), since the previous analysis has shown that the solution w given by (3.17) with (3.23) decays to zero at infinity. If $f \in \tilde{L}^2(\mathbb{R})$ does not satisfy (3.24), then no such solution $w \in \operatorname{dom}(B_0)$ exists. Hence, there exist $f \in \tilde{L}^2(\mathbb{R})$ such that for all $w \in \operatorname{dom}(B_0)$ we have $(B_0 - \mu I)w \neq f$, i.e., $\operatorname{ran}(B_0 - \mu I) \subsetneq \tilde{L}^2(\mathbb{R})$. This implies that this μ belongs to $\sigma_r(B_0)$.

In the special case $\mu = 0$, the constraint (3.24) coincides with $\langle f, \varphi \rangle = 0$. For $\mu = 0$ the unique solution (3.17) with C as in (3.18) can be rewritten as

$$(3.25) \quad w(y) = \int_{-\infty}^y \frac{\cosh(y)}{\cosh(y')} f(y') \, dy'.$$

If $\langle f, \varphi \rangle = 0$, then the solution (3.25) belongs to $H^1(\mathbb{R})$. The constraint $\langle w, \varphi \rangle = 0$, however, is satisfied only under the additional constraint

$$(3.26) \quad \int_{\mathbb{R}} \int_{-\infty}^y \operatorname{sech}(y') f(y') \, dy' \, dy = 0.$$

Therefore, for $\mu = 0$, there exists no solution $w \in \operatorname{dom}(B_0)$ to the resolvent equation (3.16) unless $f \in \tilde{L}^2(\mathbb{R})$ satisfies (3.26). This implies again that $\operatorname{ran}(B_0) \subsetneq \tilde{L}^2(\mathbb{R})$ and so $0 \in \sigma_r(B_0)$. All together we have established that $\sigma_r(B_0)$ is given by (3.14).

Finally, if $\mu = \pm 1$, one of the two exponential functions $e^{(\mu+1)y}$ and $e^{(\mu-1)y}$ in (3.17) does not decay to zero both as $y \rightarrow +\infty$ and $y \rightarrow -\infty$. Moreover, the improper integral in (3.17) does not converge for $f \in L^2(\mathbb{R})$, $f \notin L^1(\mathbb{R})$ because $e^{\pm y'} \operatorname{sech}(y') \rightarrow 2$ as $y' \rightarrow \pm\infty$. Therefore, the solution w in (3.17) does not decay to zero and does not belong to $\operatorname{dom}(B_0)$ independently on the constraint on C , and hence $(B_0 - \mu I)^{-1} : \tilde{L}^2(\mathbb{R}) \rightarrow \tilde{L}^2(\mathbb{R})$ is unbounded. We conclude that such μ belongs to $\sigma_c(B_0)$ given by (3.15). \square

Corollary 1. *The spectrum of A_0 completely covers the closed vertical strip given by*

$$(3.27) \quad \sigma(A_0) = \left\{ \lambda \in \mathbb{C} : -\frac{\pi}{6} \leq \operatorname{Re}(\lambda) \leq \frac{\pi}{6} \right\}.$$

Proof. The result follows from Lemmas 2, 3, and 4. \square

Step 4 (Justification of the truncation). In this last step, we verify that the assumptions of the abstract Theorem 1 hold for our operators. Indeed, by Lemmas 2 and 3, we have $\sigma_p(A_0) = \sigma_p(B_0) = \emptyset$. Therefore, $\rho(A) \cap \sigma_p(A_0) = \emptyset$. Moreover, Lemma 1 states that $\sigma_p(A) = \{0\}$; hence Corollary 1 implies that $\rho(A_0) \cap \sigma_p(A) = \emptyset$. Therefore, we may conclude from Theorem 1 that $\sigma(A) = \sigma(A_0)$, which together with (3.27) yields (2.5). This finishes the proof of Theorem 2.

Remark 4. We can generalize our instability result from coproduct perturbations to *subharmonic* and *localized perturbations* by analyzing the Floquet-Bloch spectrum. In particular, we find that the spectrum of A remains *invariant* with respect to the Floquet exponent k in the following decomposition:

$$v(z) = e^{ikz} p(z),$$

where $p(z + 2\pi) = p(z)$ and $k \in [-\frac{1}{2}, \frac{1}{2}]$. By setting $z = \pi \tanh(y)$, $v(z) = \cosh(y)w(y)$ as in Lemma 2, we rewrite the resolvent equation (3.16) in the following form:

$$\frac{dq}{dy} - \tanh(y)q(y) + ik\pi \operatorname{sech}^2(y)q(y) - \mu q(y) = g(y),$$

with $q(y) = e^{-ik\pi \tanh(y)}w(y)$ and $g(y) = e^{-ik\pi \tanh(y)}f(y)$. The general solution of this differential equation is obtained from (3.17) and given by

$$q(y) = \cosh(y)e^{\mu y - ik\pi \tanh(y)} \left[C + \int_0^y e^{-\mu y' + ik\pi \tanh(y')} \operatorname{sech}(y') g(y') \, dy' \right].$$

Since k is real, the analysis of this solution is exactly the same as that of (3.17) in the proof of Lemma 4. The estimates are independent of k ; therefore the spectrum of the linearized operator A remains the same when the co-periodic perturbations are replaced by subharmonic or localized perturbations.

Remark 5. If the constraint in (3.12) is dropped, one can define the differential operator $\tilde{B}_0 : H^1(\mathbb{R}) \subset L^2(\mathbb{R}) \rightarrow L^2(\mathbb{R})$, where \tilde{B}_0 has the same differential expression as B_0 in (3.10). The proofs of Lemmas 3 and 4 are extended with little modifications to show that $\sigma_p(\tilde{B}_0) = \emptyset$, $\sigma_r(\tilde{B}_0) = \sigma_r(B_0)$, and $\sigma_c(\tilde{B}_0) = \sigma_c(B_0)$. In addition, the same location of the spectrum of \tilde{B}_0 follows by Lemma 6.2.6 in [2]. Indeed, the adjoint operator $\tilde{B}_0^* : H^1(\mathbb{R}) \subset L^2(\mathbb{R}) \rightarrow L^2(\mathbb{R})$ is defined by

$$(\tilde{B}_0^* w)(y) := -\partial_y w(y) - \tanh(y)w(y), \quad y \in \mathbb{R},$$

and the exact solution of the differential equation

$$-\frac{dw}{dy} - \tanh(y)w(y) = \mu w(y)$$

is given by

$$w(y) = Ce^{-\mu y} \operatorname{sech}(y),$$

where C is arbitrary. From the decay of exponential functions, we verify directly that $\sigma_p(\tilde{B}_0^*)$ is given by (3.14) and $\sigma_c(\tilde{B}_0^*)$ is given by (3.15). However, since $\sigma_p(\tilde{B}_0) = \emptyset$, Lemma 6.2.6 in [2] implies that $\sigma_r(\tilde{B}_0^*) = \emptyset$, $\sigma_p(\tilde{B}_0^*) = \sigma_r(\tilde{B}_0)$, and $\sigma_c(\tilde{B}_0^*) = \sigma_c(\tilde{B}_0)$, which is in agreement with the location of $\sigma_r(\tilde{B}_0)$ and $\sigma_c(\tilde{B}_0)$ obtained from direct computation.

4. PROOF OF THEOREM 3

For the peaked periodic wave U_* in (1.6) in the case $p = 2$, we write explicitly

$$(4.1) \quad c_* - U_*^2(z) = \frac{1}{2}|z|(\pi - |z|), \quad z \in [-\pi, \pi].$$

The eigenvector for $0 \in \sigma_p(A)$ is given by

$$(4.2) \quad U_*'(z) = \frac{1}{\sqrt{2}}\operatorname{sign}(z), \quad z \in (-\pi, \pi).$$

We follow the same four steps as in the proof of Theorem 2. Note that now there exist two peaks of the periodic wave (1.6) on the 2π -period: one is located at $z = \pm\pi$, and the other one is located at $z = 0$. This modifies the proofs of Lemmas 1 and 2 in Steps 1 and 2, whereas Steps 3 and 4 are exactly as in the case $p = 1$.

Step 1 (Point spectrum of A). The following lemma is an adaptation of Lemma 1 for the case $p = 2$.

Lemma 5. $\sigma_p(A) = \{0\}$.

Proof. If $f \in \operatorname{dom}(A)$, then $f \in H^1(-\pi, 0) \cap H^1(0, \pi)$ so that $f \in C^0(-\pi, 0) \cap C^0(0, \pi)$ by Sobolev embedding. Bootstrapping arguments for $Af = \lambda f$ immediately yield that $f \in C^\infty(-\pi, 0) \cap C^\infty(0, \pi)$. Hence, the spectral problem $Af = \lambda f$ for $f \in \operatorname{dom}(A)$ can be differentiated once in z on $(-\pi, 0)$ and $(0, \pi)$ to yield the second-order differential equation

$$(4.3) \quad |z|(\pi - |z|)f''(z) + 2\operatorname{sign}(z)(\pi - 2|z|)f'(z) = 2\lambda f'(z), \quad z \in (-\pi, 0) \cup (0, \pi).$$

Integrating (4.3) separately for $\pm z \in (0, \pi)$ yields

$$(4.4) \quad f'(z) = \frac{g_{\pm}}{z^2(\pi - |z|)^2} \left(\frac{z}{\pi - |z|} \right)^{\pm \frac{2\lambda}{\pi}}, \quad \pm z \in (0, \pi),$$

where g_{\pm} are constants of integration. Computing the limits $z \rightarrow 0$ and $z \rightarrow \pm\pi$ similarly to the proof of Lemma 1 shows that $|z|(\pi - |z|)f'(z)$ belongs to $L^2(-\pi, 0) \cap L^2(0, \pi)$ if and only if $g_+ = g_- = 0$. In this case, $f(z) = f_{\pm}$ for $\pm z \in (0, \pi)$ with constant f_{\pm} and the zero-mass constraint $\int_{-\pi}^{\pi} f(z) dz = 0$ required for $f \in \dot{L}_{\text{per}}^2$ yields $f_{\pm} = \pm f_0$ with only one scaling constant f_0 . Hence the only solution of $Af = \lambda f$ with $f \in \text{dom}(A) \subset \dot{L}_{\text{per}}^2$ is given by $f(z) = f_0 \text{sign}(z) = \sqrt{2}f_0 U'_*(z)$ given by (4.2). Inspecting A in (2.2) with $p = 2$ shows that $(Af)(z)$ is even in z , whereas $\lambda f(z)$ is odd in z . Hence, $\lambda = 0$ is the only admissible value of λ for this solution. No other $\lambda \in \mathbb{C}$ exists such that a nonzero solution f of $Af = \lambda f$ belongs to $\text{dom}(A) \subset \dot{L}_{\text{per}}^2$. \square

Step 2 (Truncation of \mathbf{A}). By using (4.1), A_0 in (2.4) is rewritten in the explicit form

$$(4.5) \quad (A_0 v)(z) = \frac{1}{2} \partial_z [|z|(\pi - |z|)v(z)], \quad z \in (-\pi, \pi).$$

The explicit expression (4.1) in the transformation formula (1.7) for $p = 2$ yields

$$(4.6) \quad \frac{dz}{d\xi} = \frac{1}{2} |z|(\pi - |z|).$$

Both $z = \pm\pi$ and $z = 0$ are critical points of (4.6), so the interval $[-\pi, \pi]$ cannot be mapped bijectively to \mathbb{R} as in the case $p = 1$. However, we are able to map the half-intervals $[-\pi, 0]$ and $[0, \pi]$ between the two peaks separately to \mathbb{R} . These maps are given explicitly as the solutions of (4.6) by

$$(4.7) \quad z = z_+(\xi) := \frac{\pi e^{\frac{\pi\xi}{2}}}{1 + e^{\frac{\pi\xi}{2}}}$$

for $z \in [0, \pi]$, and

$$(4.8) \quad z = z_-(\xi) := -\frac{\pi}{1 + e^{\frac{\pi\xi}{2}}},$$

for $z \in [-\pi, 0]$, where the constants of integration are defined without loss of generality from the conditions $z_{\pm}(0) = \pm \frac{\pi}{2}$. The following is an adaptation of Lemma 2 when $p = 2$.

Lemma 6. *The spectral problem $A_0 v = \lambda v$ with $A_0 : \text{dom}(A_0) \subset \dot{L}_{\text{per}}^2 \rightarrow \dot{L}_{\text{per}}^2$ given by (4.5) is equivalent to the spectral problem $B_0 w = \mu w$ with*

$$(4.9) \quad \mu = \frac{4}{\pi} \lambda,$$

where $B_0 : \text{dom}(B_0) \subset \tilde{L}^2(\mathbb{R}) \rightarrow \tilde{L}^2(\mathbb{R})$ is the same linear operator as is given in (3.10) with the domain (3.11).

Proof. First, we consider the problem on the half-interval $[0, \pi]$. By setting $y := \frac{\pi}{4}\xi$ and $v(z_+) = \cosh(y)w_+(y)$, we obtain by the substitution rule and using (4.7) that

$$\int_0^{\pi} v^2(z) dz = \frac{\pi}{2} \int_{-\infty}^{\infty} v^2(z_+) \text{sech}^2(y) dy = \frac{\pi}{2} \int_{-\infty}^{\infty} w_+^2(y) dy;$$

hence $v \in L^2(0, \pi)$ if and only if $w_+ \in L^2(\mathbb{R})$. Similarly, we verify that

$$\partial_z [z(\pi - z)v] \in L^2(0, \pi)$$

if and only if

$$\partial_y w_+ - \tanh(y)w_+ \in L^2(\mathbb{R}).$$

Next, we consider the problem on the half-interval $[-\pi, 0]$. By setting $v(z_-) = \cosh(y)w_-(y)$ and using (4.8), we obtain by the same computations that $v \in L^2(-\pi, 0)$ if and only if $w_- \in L^2(\mathbb{R})$, whereas

$$\partial_z [z(\pi + z)v] \in L^2(-\pi, 0)$$

if and only if

$$\partial_y w_- - \tanh(y)w_- \in L^2(\mathbb{R}).$$

The zero-mean constraint in \dot{L}^2_{per} is transformed as follows:

$$\begin{aligned} 0 &= \int_{-\pi}^{\pi} v(z)dz = \frac{\pi}{2} \int_{\mathbb{R}} [v(z_-) + v(z_+)] \operatorname{sech}^2(y)dy \\ &= \frac{\pi}{2} \int_{\mathbb{R}} [w_-(y) + w_+(y)] \operatorname{sech}(y)dy. \end{aligned}$$

Therefore, $v \in \dot{L}^2_{\text{per}}$ if and only if $w \in \tilde{L}^2(\mathbb{R})$, where $w := w_+ + w_-$ and $\tilde{L}^2(\mathbb{R})$ is defined by (3.12). In view of (3.13) we find that $B_0w \in \tilde{L}^2(\mathbb{R})$ for $w = w_+ + w_- \in H^1(\mathbb{R})$. Considering the differential equation $A_0v = \lambda v$ on the half-intervals $[-\pi, 0]$ and $[0, \pi]$, we use the relations $v(z_{\pm}) = \cosh(y)w_{\pm}(y)$, the chain rule, and the transformation formula (4.9) to obtain the equation $B_0w_{\pm} = \mu w_{\pm}$, where the differential expression for B_0 is given by (3.10). By the linear superposition principle, $w \in \operatorname{dom}(B_0) \subset \tilde{L}^2(\mathbb{R})$ defined by (3.11) satisfies the same equation $B_0w = \mu w$ as w_+ and w_- . Hence, the spectral problems for A_0 and B_0 are equivalent to each other, and the spectral parameters λ and μ are related by the transformation formula (4.9). □

Step 3 (Spectrum of the truncated operator \mathbf{A}_0). Since the operator B_0 in Lemma 6 is identical with the one in Lemma 2, the results of Lemmas 3 and 4 apply directly to the case $p = 2$ and give the following result.

Corollary 2. *The spectrum of A_0 completely covers the closed vertical strip given by*

$$(4.10) \quad \sigma(A_0) = \left\{ \lambda \in \mathbb{C} : -\frac{\pi}{4} \leq \operatorname{Re}(\lambda) \leq \frac{\pi}{4} \right\}.$$

Step 4 (Justification of the truncation). In this last step, we verify that the assumptions of the abstract Theorem 1 hold also in the case $p = 2$. Since $\sigma_p(A_0) = \emptyset$, $\rho(A) \cap \sigma_p(A_0) = \emptyset$. Furthermore, Lemma 5 states that $\sigma_p(A) = \{0\}$; hence Corollary 2 implies that $\rho(A_0) \cap \sigma_p(A) = \emptyset$. Therefore, we may conclude from Theorem 1 that $\sigma(A) = \sigma(A_0)$, which together with (4.10) yields (2.6). This finishes the proof of Theorem 3.

APPENDIX: PROOF OF THEOREM 1

Assume that $\lambda \in \sigma(A_0)$ but $\lambda \in \rho(A)$. Hence, for every $f \in \text{dom}(A)$, we can write

$$(A.1) \quad f = (A - \lambda I)^{-1}(K + A_0 - \lambda I)f,$$

where $(A - \lambda I)^{-1} : X \rightarrow X$ is a bounded operator. The operator $(A - \lambda I)^{-1}K : X \rightarrow X$ is compact as a composition of bounded and compact operators. Therefore, the spectrum of $I - (A - \lambda I)^{-1}K$ in X consists of eigenvalues accumulating at 1. Therefore, the Fredholm alternative holds: (i) either this operator is invertible for this λ with a bounded inverse or (ii) there exists $f_0 \in \text{dom}(A)$, $f_0 \neq 0$, such that $f_0 = (A - \lambda I)^{-1}Kf_0$.

In the case (i), we can rewrite (A.1) for every $f \in \text{dom}(A)$ in the form

$$(A.2) \quad f = (I - (A - \lambda I)^{-1}K)^{-1}(A - \lambda I)^{-1}(A_0 - \lambda I)f,$$

from which we obtain a contradiction against the assumption $\lambda \in \sigma(A_0)$. Indeed, if $\lambda \in \sigma_p(A_0)$, then there exists $f_0 \in \text{dom}(A_0)$, $f_0 \neq 0$, such that $(A_0 - \lambda I)f_0 = 0$, in which case equation (A.2) yields that $f_0 = 0$, a contradiction. On the other hand, if $\lambda \in \sigma_r(A_0)$, then there exists $g_0 \in X$ such that $g_0 \notin \text{ran}(A_0 - \lambda I)$. This is in contradiction with (A.2) since for every $g_0 \in X$, there exists a unique $f_0 \in \text{dom}(A)$ such that

$$(A - \lambda I)(I - (A - \lambda I)^{-1}K)f_0 = g_0 = (A_0 - \lambda I)f_0.$$

Finally, if $\lambda \in \sigma_c(A_0)$, then for $f \in \text{dom}(A_0)$ we let $g := (A_0 - \lambda I)f \in X$ and obtain from (A.2) that

$$(A.3) \quad \|f\|_X = \|(I - (A - \lambda I)^{-1}K)^{-1}(A - \lambda I)^{-1}g\|_X \leq C\|g\|_X$$

for some $C > 0$. Since $\lambda \in \sigma_c(A_0)$, we have $\text{ran}(A_0 - \lambda I) = X$ for this λ , and since $f \in \text{dom}(A_0)$ is arbitrary, the bound (A.3) implies that for every $g \in X$,

$$\|(A_0 - \lambda I)^{-1}g\|_X \leq C\|g\|_X,$$

in contradiction with the assumption $\lambda \in \sigma_c(A_0)$.

In the case (ii), there exists $f_0 \in \text{dom}(A)$, $f_0 \neq 0$, such that $f_0 = (A - \lambda I)^{-1}Kf_0$, and hence we can rewrite (A.1) for this f_0 as

$$(A - \lambda I)^{-1}(A_0 - \lambda I)f_0 = 0.$$

Therefore, we have $(A_0 - \lambda I)f_0 = 0$, and hence $\lambda \in \sigma_p(A_0)$, in contradiction with the assumption that the intersection $\sigma_p(A_0) \cap \rho(A)$ is empty.

Thus, if $\lambda \in \sigma(A_0)$, then $\lambda \in \sigma(A)$. Since $A_0 - A = -K$ and the previous argument does not depend on the sign of K , the reverse statement is true. Hence, $\sigma(A) = \sigma(A_0)$. \square

ACKNOWLEDGMENT

Both authors thank the referee for useful remarks.

REFERENCES

- [1] G. Bruell and R. N. Dhara, *Waves of maximal height for a class of nonlocal equations with homogeneous symbol*, arXiv:1810.00248v1, 2018.
- [2] Theo Bühler and Dietmar A. Salamon, *Functional analysis*, Graduate Studies in Mathematics, vol. 191, American Mathematical Society, Providence, RI, 2018. MR3823238

- [3] Carmen Chicone and Yuri Latushkin, *Evolution semigroups in dynamical systems and differential equations*, Mathematical Surveys and Monographs, vol. 70, American Mathematical Society, Providence, RI, 1999. MR1707332
- [4] Mats Ehrnström, Mathew A. Johnson, and Kyle M. Claassen, *Existence of a highest wave in a fully dispersive two-way shallow water model*, Arch. Ration. Mech. Anal. **231** (2019), no. 3, 1635–1673, DOI 10.1007/s00205-018-1306-5. MR3902471
- [5] E. R. Johnson and R. H. J. Grimshaw, *The modified reduced Ostrovsky equation: integrability and breaking*, Phys. Rev. E **88** (2014), 021201(R) (5 pages).
- [6] Edward R. Johnson and Dmitry E. Pelinovsky, *Orbital stability of periodic waves in the class of reduced Ostrovsky equations*, J. Differential Equations **261** (2016), no. 6, 3268–3304, DOI 10.1016/j.jde.2016.05.026. MR3527630
- [7] Anna Geyer and Dmitry E. Pelinovsky, *Spectral stability of periodic waves in the generalized reduced Ostrovsky equation*, Lett. Math. Phys. **107** (2017), no. 7, 1293–1314, DOI 10.1007/s11005-017-0941-3. MR3685174
- [8] Anna Geyer and D. Pelinovsky, *Linear instability and uniqueness of the peaked periodic wave in the reduced Ostrovsky equation*, SIAM J. Math. Anal. **51** (2019), no. 2, 1188–1208, DOI 10.1137/18M117978X. MR3936896
- [9] R. Grimshaw, *Evolution equations for weakly nonlinear, long internal waves in a rotating fluid*, Stud. Appl. Math. **73** (1985), no. 1, 1–33, DOI 10.1002/sapm19857311. MR797556
- [10] R. H. J. Grimshaw, Karl Helfrich, and E. R. Johnson, *The reduced Ostrovsky equation: integrability and breaking*, Stud. Appl. Math. **129** (2012), no. 4, 414–436, DOI 10.1111/j.1467-9590.2012.00560.x. MR2993126
- [11] R. H. J. Grimshaw, L. A. Ostrovsky, V. I. Shrira, and Yu. A. Stepanyants, *Long nonlinear surface and internal gravity waves in a rotating ocean*, Surv. Geophys. **19** (1998), 289–338.
- [12] Roger Grimshaw and Dmitry Pelinovsky, *Global existence of small-norm solutions in the reduced Ostrovsky equation*, Discrete Contin. Dyn. Syst. **34** (2014), no. 2, 557–566, DOI 10.3934/dcds.2014.34.557. MR3094592
- [13] Sevdzhan Hakkaev, Milena Stanislavova, and Atanas Stefanov, *Periodic traveling waves of the regularized short pulse and Ostrovsky equations: existence and stability*, SIAM J. Math. Anal. **49** (2017), no. 1, 674–698, DOI 10.1137/15M1037901. MR3614681
- [14] S. Hakkaev, M. Stanislavova, and A. Stefanov, *Spectral stability for classical periodic waves of the Ostrovsky and short pulse models*, Stud. Appl. Math. **139** (2017), no. 3, 405–433, DOI 10.1111/sapm.12166. MR3708844
- [15] Aleksey Kostenko and Noema Nicolussi, *On the Hamiltonian-Krein index for a non-self-adjoint spectral problem*, Proc. Amer. Math. Soc. **146** (2018), no. 9, 3907–3921, DOI 10.1090/proc/14048. MR3825844
- [16] Yue Liu, Dmitry Pelinovsky, and Anton Sakovich, *Wave breaking in the Ostrovsky-Hunter equation*, SIAM J. Math. Anal. **42** (2010), no. 5, 1967–1985, DOI 10.1137/09075799X. MR2684307
- [17] S. P. Nikitenkova, Yu. A. Stepanyants, and L. M. Chikhladze, *Solutions of a modified Ostrovskii equation with a cubic nonlinearity* (Russian, with Russian summary), Prikl. Mat. Mekh. **64** (2000), no. 2, 276–284, DOI 10.1016/S0021-8928(00)00048-4; English transl., J. Appl. Math. Mech. **64** (2000), no. 2, 267–274. MR1773718
- [18] L. A. Ostrovsky, *Nonlinear internal waves in a rotating ocean*, Okeanologia **18** (1978), 181–191.
- [19] L. Ostrovsky, E. Pelinovsky, V. Shrira, and Y. Stepanyants, *Beyond the KdV: post-explosion development*, Chaos **25** (2015), no. 9, 097620, 13, DOI 10.1063/1.4927448. MR3396218
- [20] Roman Shvidkoy and Yuri Latushkin, *The essential spectrum of the linearized 2D Euler operator is a vertical band*, Advances in differential equations and mathematical physics (Birmingham, AL, 2002), Contemp. Math., vol. 327, Amer. Math. Soc., Providence, RI, 2003, pp. 299–304, DOI 10.1090/conm/327/05822. MR1991549
- [21] Atanas Stefanov, Yannan Shen, and P. G. Kevrekidis, *Well-posedness and small data scattering for the generalized Ostrovsky equation*, J. Differential Equations **249** (2010), no. 10, 2600–2617, DOI 10.1016/j.jde.2010.05.015. MR2718712
- [22] Milena Stanislavova and Atanas Stefanov, *On the spectral problem $\mathcal{L}u = \lambda u'$ and applications*, Comm. Math. Phys. **343** (2016), no. 2, 361–391, DOI 10.1007/s00220-015-2542-2. MR3477342

DELFT INSTITUTE OF APPLIED MATHEMATICS, FACULTY ELECTRICAL ENGINEERING, MATHEMATICS AND COMPUTER SCIENCE, DELFT UNIVERSITY OF TECHNOLOGY, VAN MOURIK BROEKMANWEG 6, 2628 XE DELFT, THE NETHERLANDS

Email address: A.Geyer@tudelft.nl

DEPARTMENT OF MATHEMATICS AND STATISTICS, MCMASTER UNIVERSITY, HAMILTON, ONTARIO, CANADA, L8S 4K1; AND DEPARTMENT OF APPLIED MATHEMATICS, NIZHNY NOVGOROD STATE TECHNICAL UNIVERSITY, 24 MININ STREET, 603950 NIZHNY NOVGOROD, RUSSIA

Email address: dmpeli@math.mcmaster.ca